

Magnetic reconnection: flares and coronal heating in Active Galactic Nuclei

T. Di Matteo

Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA

1 February 2008

ABSTRACT

A magnetically-structured accretion disk corona, generated by buoyancy instability in the disk, can account for observations of flare-like events in Active Galactic Nuclei. We examine how Petschek magnetic reconnection, associated with MHD turbulence, can result in a violent release of energy and heat the magnetically closed regions of the corona up to canonical X-ray emitting temperatures. X-ray magnetic flares, the after effect of the energy released in slow shocks, can account for the bulk of the X-ray luminosity from Seyfert galaxies and consistently explain the observed short-timescale variability.

Key words: - MHD - magnetic fields: flares - galaxies: active - accretion discs

1 INTRODUCTION

Active Galactic Nuclei (AGN) are believed to be powered by accretion. UV and X-ray observations of AGN, particularly in Seyfert 1 galaxies, indicate that the gravitational binding energy of a massive black-hole is dissipated partly in a cold accretion disk and partly in a hot, tenuous corona above it. Thermal Comptonization of soft UV-radiation in the corona leads to the production of a hard X-ray continuum, some of which is reprocessed by the cold, dense disk (Haardt & Maraschi 1991, 1993) giving rise to the observed reflexion features in the spectra. Recent work, both on the observational and theoretical side, (Sincell & Krolik 1997, Nandra 1997, Stern et al. 1995) indicates that the UV continuum in AGN may be mostly produced by reradiation of energy absorbed from X-rays irradiating the accretion disk. Also, in order to explain the different ratios of X-ray and UV luminosities in different objects it has been suggested that the corona consists of localized active regions (e.g. Haardt, Maraschi & Ghisellini 1994). Popular models for the production of the X-rays, therefore, suppose that a large part of the disk's dissipation takes place in a small amount of mass in the hot corona. Variability timescales of the order of few hours observed in AGN also put upper limits on the size of the emitting regions, and imply that enormous amounts of energy need to be released on very short timescales. It is therefore important to understand what processes can efficiently channel significant amounts of energy into the hot coronal medium and the way in which energy can be released in localized flare-type events.

In a differentially rotating disk the evolution of a magnetically structured corona is very hard to suppress. The

strong magnetic fields, continuously generated by the dynamo action in an accretion disk, are strongly buoyant and are forced to invade the region sandwiching the disk itself. Once outside the disk, the magnetic flux tubes can reconnect efficiently and dissipate part of the accretion energy in localized active flares. Buoyancy, therefore, constitutes a mechanism that channels part of the energy released in the accretion process directly into the corona outside the disk. Magnetic reconnection can be responsible for the rapid dissipation of magnetic energy through a field-aligned electric potential in thin current sheets.

We, here, describe how the observationally required large heating rates per unit mass, in the optically thin gas of a hot coronal region, can be accounted for by an ion-acoustic instability in the context of slow shocks associated with Petschek type reconnection and give rise to flare-like events. We estimate that such a process may be responsible for the heating of the coronal plasma in the active regions of an AGN to a level where it emits X-rays at a temperature of $\sim 10^9$ K even in the presence of Inverse Compton and synchrotron cooling processes. Magnetic buoyancy, which drives magnetic flux out of the disk, naturally decreases the plasma density to the point where the active region can reconnect very efficiently. X-ray observations imply that the coronal plasma is very tenuous with a density much lower than that of the underlying disk. We show that the energetics of such flares are consistent with the X-ray luminosities of typical Seyfert galaxies. The onset of magnetic flares through slow shocks associated with Petschek reconnection, should occur over timescales comparable to those required to explain X-ray variability observed in AGN.

This picture is also, by analogy, validated by X-ray ob-

servations of the solar corona. Recent *Yohkoh* observations (Tsuneta 1996, Yokoyama & Shibata 1995 and references therein) of solar flares show clear evidence for magnetic reconnection taking place in magnetically confined loop-like volumes. There is evidence that reconnection serves as a highly efficient engine for converting magnetic energy into kinetic and thermal energy with standing slow shocks. Moreover, it is now well established that in the Sun active regions are formed from the emergence of magnetic flux. It has been shown (Shibata et al. 1989) that magnetic reconnection can be driven self-consistently by the magnetic buoyancy instability (the Parker instability) between emerging flux and the overlying coronal magnetic field.

2 EMERGING MAGNETIC FLUX AND RECONNECTION IN AGN CORONAE

Magnetic fields have long been considered an important element in the dynamics of accretion disks in particular as a mechanism for supplying the internal stresses required for efficient angular momentum transport (Coroniti 1979, Stella & Rosner 1984, Balbus & Hawley 1991). Strong fields in the accretion disk are continuously generated because the shearing and the turbulent disk flow serve as an effective dynamo that can rapidly strengthen any seed field. Field amplification is limited by non-linear effects: as a consequence of buoyancy (Parker instability) strong fields will rapidly emerge from the disk leading to an accretion disk corona consisting of many magnetic loops (Galeev Rosner & Viana 1979). Part of the accretion energy is therefore released into a corona in the form of magnetic energy. Either by internal shear or by encountering already present magnetic flux tubes, reconnection takes place and rapid magnetic energy dissipation can occur through the formation of a slow shock in a thin current sheet.

Not all of the emerging flux will produce a flare. Simple loop flares are triggered only if sufficient flux emerges so that the current sheet occupying the interface between new and old flux reaches a critical height that depends on the emergence speed and magnetic field strength (e.g. Heyvaerts, Priest & Rust 1977 and references therein). As the sheet rises through the corona its temperature increases until it achieves a state where Joule heating is balanced by thermal conduction. At this point, the current sheet can reach the critical temperature for the onset of turbulence (creation of a slow shock). This is the main phase of the flare, the current sheet reaches a steady state, with reconnection based on a marginally turbulent resistivity (anomalous electric resistivity), with X-ray flares being the after effect of the energy release by the slow shocks.

In summary, in a differentially rotating disk gas, the Parker instability should lead to the formation of an electro-dynamically coupled corona. The onset of impulsive reconnection corresponds to the evolution of anomalous dissipation in localized regions, in a similar manner to flare events in the Sun's corona. Although the plasma regime envisaged here for the accretion disk corona is vastly different from that of the Sun, its structure is quite similar. In both cases the low- β coronal plasma is dynamically controlled by magnetic fields the footpoints of which are embedded in a buoyantly unstable, $\beta \sim 1$ plasma.

It is now well established that active regions in the Sun are formed by the emergence of magnetic flux tubes. The soft X-ray telescope aboard of *Yohkoh* has provided the opportunity to observe the interaction of emerging flux with overlying coronal magnetic fields with high spatial and temporal resolution (e.g. Tsuneta 1996). In addition, numerical simulations of magnetic reconnection, driven by magnetic buoyancy instability, have been performed by solving resistive magnetohydrodynamic equations assuming uniform magnetic field at the base of the Sun's corona. The emergence of flux sheets is reproduced by simulating the Parker instability. As the instability develops, gas slides down the expanding loop, and the evacuated loop rises as a result of the enhanced magnetic buoyancy (expansion is due to magnetic pressure force although the dynamics are controlled by the downflow, due to gravity, along the rising loop). Consequently, as long as the magnetic pressure at the top of the magnetic loop is larger than the coronal (gas and magnetic) pressure, the expansion of the magnetic loop continues. In particular detailed simulations of such instability (Shibata et al. 1989) have shown that during the expansion of the magnetic loop the rise velocity and the local Alfvén speed of the loop increases linearly with the height h above the disk whereas the density in the sheet scales as $\rho \propto h^{-4}$ (the normalizing constant is H , the scale height of the basis where the plasma β distribution initiates the instability).

In order to meet the observational constraints, the coronal plasma in an AGN needs to be very tenuous (optically thin). The optical depth deduced from X-ray observations is $\tau \approx 1$ (e.g. Poutanen 1997 – where τ is the Thomson optical depth given by $\tau = n\sigma_T R$, n is the electron number density and σ_T the Thomson cross section). Since AGN show variations in luminosity on very short time scales, R , the thickness of the emitting region, can be constrained from observations of the shortest variability timescales ($R/c \sim 200$ s) to be of the order $R \sim 6 \times 10^{12}$ cm. For τ in the given range, $n \sim 3 \times 10^{11} \text{ cm}^{-3}$. If the energy released in a flare needs to account for an impulsive event and provide the required energy dissipation rates, reconnection needs to be very efficient. Clearly, as the reconnection velocity increases ($\propto V_A$ the Alfvén speed; $V_A = B/\sqrt{4\pi n m_p}$) where B is the magnetic field strength and m_p the proton mass) the timescale for reconnection decreases and the energy dissipation rate increases. Therefore, according to magnetohydrodynamic theory, reconnection is most efficient when the current sheet density is low or equivalently when the loop has been pushed up high enough in the corona. As mentioned above, the density would naturally decrease as the magnetic tube expands and rises from an underlying dense disk (with, typically, $n_{\text{disk}} \sim 10^{16} \text{ cm}^{-3}$, $T_{\text{disk}} \sim 10^5$ K) due to the gravitational downflow caused by the magnetic buoyancy instability and reach values of the order of n as determined above from observational constraints. If we assume that a flare is triggered when the density has decreased to value $n_{in} \sim 2 \times 10^{12} \text{ cm}^{-3}$ (n_{in} is the density of the material entering the shock: a factor 5–6 greater than n as obtained from the jump condition, eqn. (6)) we can estimate the critical height at which reconnection could suddenly start in an AGN corona. According to the afore described magnetic buoyancy simulations, the loop expansion is self-similar and the loop top reaches a height given by

$$h_{\text{flare}} \sim (10^{16}/2 \times 10^{12})^{0.25} H \sim 8H, \quad (1)$$

where H , in this case, is the pressure scale height of the accretion disk (in a standard Shakura & Sunyaev model $H/R \lesssim 0.1$, so that in the central regions ($R = 3 - 4R_S$), h_{flare} would correspond to about $3R_S$, where R_S is the Schwarzschild radius). This implies that a magnetically structured corona can exist at a few scale heights above the disk. This is consistent with theoretical arguments related to buoyancy: the greater shear energy in the larger magnetic flux tubes in the disk amplifies the magnetic field beyond equipartition and therefore allows only the large scale length flux ($R \gtrsim H$) to be expelled from the disk. (There are no obvious constraints from the spectra of Seyfert galaxies that can rule out the occurrence of flares at scale heights larger than H).

Given the relevance of emergence flux models in coronae, it would be appropriate to treat magnetic reconnection in a self-consistent manner as the interaction and coalescence of the rising flux tubes. Such a treatment is still a very difficult task as it would require a run of densities magnetic field with h . We will therefore consider only the current sheet in a self-consistent manner adopting values for the density interpreted from observations of AGN X-ray emission.

3 MAGNETIC RECONNECTION AND CORONAL HEATING

In this section we describe the dynamics of the formation of localized field-aligned electric fields in the corona of an AGN within the context of a simple analytical model.

The major problem faced by any mechanism which attempts to dissipate the stored magnetic energy is the low resistivity of the pre-flare coronal plasma, with associated long magnetic diffusion times and high magnetic Reynolds numbers ($R_m = \tau_D/\tau_A$, where τ_D is the diffusion timescale and τ_A is the Alfvén time). Since the diffusion timescales varies as the squares of the characteristic length scale, l , associated with changes of magnetic field, it is desirable to make l as small as possible. The topology of reconnection offers a way of achieving this, i.e. two sets of oppositely directed fields in close proximity – a so called neutral sheet configuration.

The current density in a neutral sheet is given by

$$j = nev = \frac{c}{4\pi} (\nabla \times \mathbf{B}) \sim \frac{cB_x}{4\pi l}. \quad (2)$$

As l becomes small, j becomes large and the Ohmic dissipation ηj^2 can become very large, where η is the magnetic resistivity. Moreover, where magnetic reconnection is required to explain various explosive phenomena, Petschek models need to be invoked. Petschek reconnection takes into account the inclusion of the effect of waves – the occurrence of a slow mode MHD shock – and is almost independent of Reynolds numbers. The development of plasma instabilities (turbulence in the shock) implies the onset of anomalous resistivity much larger than the classical resistivity. The electrons velocities are randomized by scattering on the waves and, in this manner, the dissipated energy goes mainly into heating of the resonant electrons. A well-studied example of such a mechanism (and probably most important mechanism in the neutral sheet) is the ion acoustic instability that

sets in when the electron drift velocity surpasses the ion-acoustic wave velocity (e.g. Coroniti & Eviatar 1977). This provides fast reconnection rates leading to explosive outbursts. Simulations (Yokoyama & Shibata 1994) have shown that for magnetic reconnection driven by the Parker instability anomalous instability indeed leads to (fast) Petschek-type reconnection.

In the following section we describe the magnetic structure and dynamics of the reconnection region of the flare. We employ the theory of Petschek reconnection (X-point with slow shock structure) which appears to be consistent with the detailed observations of the Sun's coronal flares.

3.1 Slow standing shock

We here review the basic set of equations which are useful for our calculations. The inflow velocity toward the X-point and the slow shocks is V , and the outflows have the Alfvén speed in both the Sweet–Parker or in the Petschek theories (momentum balance). The latter theory also predicts that the acute half angle of the slow shock, $\alpha \approx 1 - 2^\circ$ (this is supported by observations). Mass conservation then gives

$$n_{\text{in}} V L \cos \alpha = n_{\text{out}} V_A L \sin \alpha, \quad (3)$$

where L is the length of the region of close, oppositely directed magnetic field lines (the length of the slow shock region). The suffixes n_{in} and n_{out} indicate the inflow and the outflow respectively. We thus obtain

$$V = V_A \tan \alpha \frac{n_{\text{out}}}{n_{\text{in}}}. \quad (4)$$

We next examine the standing slow shock attached to the X-point. It is usually assumed that the slow shock is switch-off, and that the inflow speed $V \lesssim V_A$. The momentum flux across the shock must be continuous giving

$$p_{\text{in}} + \frac{B_{\text{in}}^2}{8\pi} = p_{\text{out}}, \quad (5)$$

where p_{in} and p_{out} are the inflow and outflow gas pressures respectively. Since the thermal conduction along the reconnected field lines is high, we expect the slow shock to be isothermal within the thermal conduction front (across the shock region). Therefore the jump condition for the plasma density is given from eqn. (5)

$$\frac{n_{\text{in}}}{n_{\text{out}}} = 1 + \frac{1}{\beta_{\text{in}}}, \quad (6)$$

where the plasma β is given by $\beta = p_{\text{in}}/(B_{\text{in}}^2/8\pi)$, which for Petschek reconnection is typically small (of the order of $0.2 - 0.3$, giving a density jump of $\sim 5 - 6$).

In order to set the typical magnetic field strength, B , in the rising loop we use the condition for the onset of magnetic buoyancy in the disk. If the shear stresses acting on a flux cell generate a magnetic pressure which exceeds the gas pressure, upward buoyancy forces will rapidly remove the flux tubes from the disk (Rosner & Vaiana 1979 and Coroniti 1981). The field strength in the loop is set by the condition that $\beta \sim 1$ or

$$\frac{B^2}{8\pi} \approx n_{\text{disk}} k T_{\text{disk}}. \quad (7)$$

In order to maintain pressure balance, magnetic flux tubes with such strong magnetic fields will contain less plasma

than their ambient surroundings; therefore, they become subject to buoyancy and will emerge from the disk as described in the previous sections. For a typical AGN, the disk temperature is of the order of $T_{\text{disk}} \sim 10^5$ K and its number density $n_{\text{disk}} \sim 10^{16} \text{ cm}^{-3}$. B is therefore $\sim 2 \times 10^3$ Gauss. With this value of B , $V_A = B/\sqrt{(4\pi n_{\text{out}} m_p)} \sim 10^9 \text{ cm s}^{-1}$ where $n_{\text{out}} = n \sim 3 \times 10^{11} \text{ cm}^{-3}$, $V = 1/10 V_A$ and from continuity, $l/L \approx V/V_A = 1/10$. We take the dimension l to be set by shortest variability timescale i.e. $l = R = 6 \times 10^{12} \sim 2R_s$ where R_s is the Schwarzschild radius for a mass $M = 10^7 M_\odot$, typical of a Seyfert.

3.2 Energetics

We next examine the energetics of a reconnection site when the main phase of a flare is triggered i.e. when turbulence is created with an associated MHD slow shock. Our aim is to obtain order of magnitude estimates of the different processes which contribute to the heating and cooling of the current sheet and to derive an estimate of the energy released in a flaring corona in an AGN.

The plasmas on the reconnected field lines are significantly heated by the two standing slow shocks. In addition to the heating, radiative losses, especially due to Inverse Compton and synchrotron processes, become important in an AGN environment and will also be taken into account.

We consider one single sheared coronal loop, i.e. a magnetic flux tube with a current flowing due to the magnetic shear. We assume that the energy produced in the reconnection site goes into slow MHD shock heating as well as energy of the bulk flow. This assumption is good for the Petschek reconnection model, where the slow shock angle is small and the outflow plasma β_{out} is high. Energy balance therefore gives

$$\frac{B_{\text{in}}^2}{4\pi} V L = \frac{1}{2} (m_p n_{\text{out}} V_A^2 + 5 n_{\text{out}} k T) V_A l + \kappa_0 \frac{T^{7/2}}{L} l + E_r. \quad (8)$$

i.e. the energy dissipation rate has to balance with the inflow magnetic energy. The term on the left hand side of equation (8) represents the magnetic enthalpy flux which coincides with $(1/4)$ of Joule heating of the current sheet ($\eta j^2 4Ll$) (eqn. 2). The first two terms on the right hand side are the kinetic energy flux and the enthalpy flux of the outflow. The third term on the right hand side is the heat flux and represents an estimate of the heat produced in the slow MHD shock from conduction losses. The term $\kappa_0 T^{5/2} (T/L)$ is the Spitzer thermal conductivity ($\kappa_0 = 1.79 \times 10^{-5} / \ln \Lambda$) and T is the temperature of the slow MHD shock. The forth term represents the relevant radiative energy losses. The electrons in the coronal loop are cooled through inverse Compton scattering and synchrotron radiation. Hence

$$\begin{aligned} E_r/4Ll &= q_{\text{Comp}} + q_{\text{synch}} \\ &= 4 \left(\frac{kT}{m_e c^2} \right) \sigma_T n_{\text{out}} c U_{\text{rad}} + \frac{2\pi}{3} \frac{kT}{c^2} \frac{\nu_c^3}{l}, \end{aligned} \quad (9)$$

where σ_T is the Thomson cross section and U_{rad} is the energy density of the isotropic soft photon bath. The seed photons are partly created locally by thermal synchrotron processes and partly created in the underlying disk. The photons from the disk are in turn due to both local energy dissipation and to re-radiation of hard X-rays created in the active




Figure 1. The temperature solution for a current sheet producing a flare in an AGN for different values of the optical depth τ . The dashed and solid lines represent the temperature solution for $U_{\text{rad}} = (B^2/8\pi)V_A/c$. The thinner lines (dotted and dot-dashed) show the effect of considering greater Inverse Compton losses, i.e. $U_{\text{rad}}(\text{max}) = (B^2/8\pi)$. The solid and dot-dashed lines take into account the effects of the slow MHD shock on the conduction term (anomalous conduction F_{an} as opposed to E_H).

coronal loops. Recent work (Nandra 1996; Sincell & Krolik 1997) indicates that most of the quasi-thermal emission from the disk is dominated by reradiation of the energy absorbed from the X-rays irradiating the disk. This implies that U_{rad} can not exceed $U_{\text{mag}} = B^2/8\pi$ (i.e. the total energy density stored in the magnetic structure) and depends on how efficiently magnetic field is dissipated. A reasonable photon energy density is then given by $U_{\text{rad}} = (B^2/8\pi)V_A/c$ where c is the light speed (see also Di Matteo, Blackman & Fabian 1997). ν_c is the self-absorbed synchrotron frequency is obtained by equating the Rayleigh-Jeans emission to the synchrotron emission (Zdziarski 1985).

The classical thermal conductivity, $E_H = \kappa_0 T^{7/2}/L$ only represents a lower limit to the heating that goes into the slow shock region. When the anomalous resistivity (with a slow MHD shock) develops the conductivity reduced (i.e. it saturates) with respect to its classical value. Taking plasma turbulence to be the source of energy release in the flare, the field-aligned thermal flux becomes anomalous and can be approximated by (Manheimer 1977; see also Somov 1992)

$$F_{\text{an}} = \frac{n_{\text{in}} (kT)^{3/2}}{4m_e^{1/2}}, \quad (10)$$

where the simple analytical function F_{an} is obtained under the assumption that a current sheet can reach an equilibrium state specified by saturated ion-acoustic turbulence. We can now solve equation (8) for both F_{an} and E_H . With B and all the relations between velocities and densities for the outflow and the inflow derived as in the previous section, we can solve for the temperature of a flare in an AGN corona, for a range of τ (densities). The resulting temperature solutions with the two respective conduction terms is shown in Fig 1. We show temperature solutions for $U_{\text{rad}} = (B^2/8\pi)V_A/c$ and for the extreme value, $U_{\text{rad}}(\text{max}) = B^2/8\pi$ (Inverse Comp-

ton losses usually dominate synchrotron ones). This clearly illustrates that temperatures of the order of a few $\times 10^9$ K can be easily achieved, even in the presence of intense radiation fields, implying that magnetic reconnection can indeed provide for the heating of the plasma to a level where it emits X-rays. Observations of spectral cut-offs in AGN directly determine the temperature of the X-ray emitting gas to be of this order (e.g. see Poutanen et al. 1997). It is worth stressing that because of the strong dependence of the shock heating term on T , the temperature solution, $T \sim 10^9$ K is very stable and only weakly dependent on changes in current sheet dimensions or B .

Note that for $\tau > 1$ (and corresponding density) the conduction cooling term acts very efficiently and the temperature of the flare becomes $< 10^9$ K. Flares with such τ would only contribute to emission in soft X-ray band and, given the strong dependence on T of the conduction flux, they would be energetically negligible. This is in accord with interpretation of X-ray observations which imply $\tau \lesssim 1$, (also discussed in the context of coronal magnetic flares by Nayakshin & Melia 1997).

The magnetic energy is converted into plasma heating in the slow shock regions to give the temperature just determined. It is important to check whether the total energy production rate for flares, as estimated from the amount of energy going into the shock (i.e. to the conduction flux), is consistent with the energy output of a typical AGN. For $T \sim 2 - 3 \times 10^9$ K, the energy of one flare is then given by

$$E_{\text{flare}} = F_{\text{an}}(4Ll) \sim 5 \times 10^{41} \text{ erg s}^{-1}. \quad (11)$$

This implies that we only require $N \sim 10 - 20$ flares to be active at any given time in order to account for typical Seyferts luminosity – of the order of $10^{42-43} \text{ erg s}^{-1}$ (actually Eqn. (11) only gives a lower limit to the total energy going into the shock and into particle heating: energy can also be tapped from internal and kinetic energy terms). Therefore, the energy produced in slow MHD shocks is completely consistent with the X-ray energy output of an AGN.

We have shown, by means of order of magnitude estimates, that the slow shock region in the current sheet of an AGN corona is heating-dominated and can give rise to a flare. In magnetically structured coronae in AGN or Galactic black hole candidates, the hard X-rays are produced by Inverse Compton scattering on 10^9 K electrons, so clearly the energy flow into the electrons has, in some way, to be guaranteed. Assuming that the current sheet becomes turbulent implies that dissipation will take place in a spectrum of length-scales, from a dominant scale down to the smallest scales, and, therefore down to the shortest possible timescales. On average, the timescale associated with the flaring of such a magnetic loop is, in accordance with Petschek models completely consistent with the typical variability timescales of Seyferts. Petschek (1964) in fact proposed that the inclusion of the effect of waves enables reconnection to be as short as a few Alfvén times; $\tau_A = l/V_A \sim 100$ a few thousand seconds depending on B and l and almost independent of the Reynolds number.

4 SUMMARY AND DISCUSSION

We have presented a simple model that takes into account the impulsive dissipation of magnetic energy during coronal flares in AGN. According to standard models, the X-rays are produced by Inverse Compton scattering of lower energy photons on energetic electrons. We show that the energy flow to the electrons that is needed for this process is guaranteed by the release of coronal magnetic energy. The onset of an ion-acoustic instability associated with slow MHD shocks and Petschek reconnection, heats the flare plasma to X-ray temperatures $\sim 10^9$ K, as required by observations. The luminosity produced in a magnetically structured, flaring corona is completely consistent with typical power outputs of AGN if at least $N \sim 10$ flares are triggered at any given time. The plasma reaches low enough densities, (high Alfvén speeds), when reconnection is driven by the Parker instability, such that flares are typically triggered at $h \sim 8H$ above the accretion disk (coronal structures can have scale-heights greater than the disk). Energetic flare-type events naturally explain the observations of short timescale variability in AGN.

The buoyantly unstable magnetic flux tubes, once outside the disk, rise through the coronal atmosphere at their local Alfvén speed. The timescale for them to rise up to a few scale heights therefore is much shorter than the disk dynamical timescale (and less than the shear timescale in the disk $\sim 2/3\Omega$). This implies that tubes will not be disrupted even when flares are triggered a few scale-heights above the disk.

An important problem faced by any model of AGN coronae, is to determine the population of non-thermal particles produced by the slow MHD shock and the effects of the impulsive electric field. According to theoretical arguments for the kind of electric fields of interest, the turbulence is characterized by an highly anisotropic distribution of ion-acoustic waves. In these circumstances it has been shown (Heyvaerts et al. 1977) that electron heating occurs with practically no acceleration. Because electrons travel much faster than ion-acoustic waves, the resonance occurs mainly with waves propagating normal to the particle velocity: this leads to angular diffusion with little change in particle energy. This implies that electrons can be characterized by a mean increase of energy or ‘temperature’. In this sense we can treat the hot current sheet as a ‘thermal’ source of electrons as required by the observations of spectral cut-offs in the X-ray spectra of AGNs. Note, though, that the presence of an electric field and plasma turbulence in the sheet would inevitably cause acceleration of charged particles to a certain degree. This is particularly relevant in the case of Galactic black hole candidates where recent high energy X-ray observations require the presence of non-thermal electron tails (e.g. Poutanen & Coppi 1988). Further investigations are needed but they are beyond the scope of this paper.

The geometry of reconnection described here does not provide the only viable way for the process to occur. In an accretion disk corona, the closed magnetic loops will get twisted by the rotation of the accretion disk. As the twist accumulates, the magnetic loops expand and finally approach the open field configuration. A current sheet is formed inside the expanding loops and, in the presence of resistivity, magnetic reconnection will take place. Open field lines an-

chored to the disk, may be the region where winds or jets blow from the disk; these could play an important role in the dynamics of the accretion disk itself. The luminosity of flares will then vary with respect to the different values of B in different morphologies in which reconnection takes place, and with the size and the total number of flares. All of these factors can give rise different covering fractions of the X-ray emitting regions which can result in very different variability timescales as observed in AGN. Finally, magnetic buoyancy is not strictly necessary in order to maintain a certain level of activity. Once a flare has been triggered, reconnection can be maintained by positive feedback (the fast outflow due to reconnection rarefies the reconnection region, thinning the current channel in this way maintaining the anomalous resistivity at the neutral sheet).

ACKNOWLEDGEMENTS

I acknowledge PPARC and Trinity College for financial support. I thank my supervisor A.C Fabian, E. Blackman, M.J. Rees and in particular Jean Heyvaerts for discussions.

REFERENCES

- Balbus S.A., Hawley J.F., 1991, ApJ, 376, 214
 Coroniti, F.V. Eviatar, 1977, ApJS, 33, 189
 Di Matteo T., Blackman E.G., Fabian A.C., 1997 MNRAS, 291, L23
 Galeev A.A., Rosner R., Vaiana G.S., 1979, ApJ, 229, 318
 Haardt F., Maraschi L., ApJ, 1993, 413, 507
 Haardt F., Maraschi L., Ghisellini G., 1994, ApJL, 432, L92
 Heyvaerts J., Priest E.R., Rust D.M., 1977, ApJ, 216, 123
 Manheimer W. M., 1977, Phys. Fluids, 20, 265
 Nayakshin S., Melia F., 1997a, 1997, ApJ, 490, L13
 Nayakshin S., Melia F., 1997b submitted to ApJL
 Nandra K., 1997, in IAU Colloq. 159, Emission Lines in Active Galactic Nuclei: New Methods and Techniques, ed. B. Peterson, F.Z., Cheng & A.S. Wilson
 Parker, E.N. *Cosmical Magnetic Fields*, (Oxford: Clarendon Press, 1979)
 Petschek, H.E., 1964, AAS-NASA Symp. on Phys. of Solar Flares, NASA SP-50, 425
 Priest, E.R. & Forbes, T.G., 1986, J. Geophys. Res. 91, 5579.
 Poutanen J., Svensson R., Stern B., 1997, Proceedings of 2nd INTEGRAL L Workshop, StMalo, France, p.401
 Poutanen J., Coppi P., 1998, to appear in Physica Scripta in the proceedings of the Nordic Symposium "Exploiting Spectrum-X-Gamma and INTEGRAL", Copenhagen
 Shibata, K., Tajima, T., Steinolfson R.S., & Matsumoto, R., 1989, ApJ, 345, 584
 Sincell M.W., Krolik J.H., 1997, ApJ, 476, 605
 Somov B.V., 1992, *Physical Processes in Solar Flares*, Kluwer Academic Publishers, Dordrecht
 Stella L., Rosner R., 1984, ApJ, 277, 312
 Stern B.E., Poutanen J., Svensson R., 1995, 449, L13
 Tsuneta S., 1996, ApJ, 456, 840
 Tsuneta S., Masuda S., Kosugi T., Sato J., 1997, 478, 787
 Yokoyama T., Shibata K., 1995, Nat., 375 42
 Yokoyama T., Shibata K., 1994, ApJ, 436, L197
 Zdziarski A.A., Johnson W.N., Done C., Smith D., McNaron-Brown K., 1995, ApJ, 438, L63